


# Up to 41%\* superior improvement in lung function vs. Spiolto Respimat (180mL vs. 128mL; $p < 0.001$ )<sup>1</sup>



\*ANORO showed superiority on the primary endpoint of trough FEV<sub>1</sub> compared to Spiolto Respimat (tiotropium/olodaterol 5/5mcg).<sup>1,6</sup> Trough FEV<sub>1</sub> improved to 180mL for ANORO (n=225) versus 128mL for Spiolto Respimat (n=224) over an 8-week treatment period in adults with symptomatic moderate COPD (mMRC  $\geq 2$  and post-bronchodilator FEV<sub>1</sub> 50–70%); difference 52mL (95% CI: 28, 77;  $p < 0.001$ ) in ITT population.<sup>1</sup>

For symptomatic patients with COPD, increasing lung function and reducing breathlessness has demonstrated a positive effect on their quality of life.<sup>7</sup> ANORO achieved superior lung function improvement and improved symptoms vs. other LAMA/LABAs within the class, helping reduce the impact of COPD on your patients' daily lives.<sup>1,3</sup>

## Trough FEV<sub>1</sub> at Week 8 in adults with moderate COPD<sup>1†</sup>



ANORO delivered dose: 55/22mcg  
Adapted from Feldman *et al* 2017

Minimum clinically important difference (MCID) defined as the minimum improvement that patients can perceive. In lung function, MCID=100 mL improvement in trough FEV<sub>1</sub> from baseline.<sup>8</sup>

<sup>†</sup>Defined as FEV<sub>1</sub>  $\leq 70\%$  to  $\geq 50\%$  predicted, mMRC  $\geq 2$ .

Anoro is subject to a Special Authority Criteria and is only funded for patients stabilised on a LAMA and who would receive additional benefit from switching to a combination product.<sup>5</sup> Anoro is well tolerated. Common adverse events include nasopharyngitis, cough and oropharyngeal pain.<sup>4</sup>

1. Feldman GJ *et al*. *Adv Ther* 2017;34(11):2518–2533. 2. Alcázar Navarrete B *et al*. *Pulm Ther* 2018;4:171–183. 3. Maltais F *et al*. *Adv Ther* 2019;36:2434–2449. 4. GlaxoSmithKline New Zealand. Anoro Ellipta Data Sheet. GSK NZ; 2020. Available at <https://medsafe.govt.nz/profs/datasheet/a/anoroelliptapowder.pdf> (Last accessed February 2021). 5. PHARMAC Special Authority for Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists. Available from <https://ieschedule.pharmac.govt.nz/ScheduleOnline.php?osq=Umeclidinium+with+vilanterol> (Last accessed February 2021). 6. Spiolto Respimat Data Sheet. Available at <https://medsafe.govt.nz/profs/datasheet/s/SpioltoRespimatinh.pdf> (Last accessed March 2021). 7. Donohue JF *et al*. *Pulm Pharmacol Ther* 2018;49:11–19. 8. Jones PW *et al*. *Am J Respir Crit Care Med* 2014; 189:250–255.

LAMA, long-acting muscarinic antagonist; LABA, long-acting  $\beta_2$ -agonist; COPD, chronic obstructive pulmonary disease; FEV<sub>1</sub>, forced expiratory volume in one second

**Anoro Ellipta** (umeclidinium bromide/vilanterol bifenatate inhaler 62.5/25mcg per inhalation) is a **Prescription Medicine**. Anoro Ellipta is indicated as a long-term maintenance bronchodilator treatment to relieve symptoms in adult patients with Chronic Obstructive Pulmonary Disease (COPD). **Anoro Ellipta is a fully funded medicine; Special Authority criteria apply. Maximum Daily Dose:** One inhalation once daily. **Contraindications:** Patients with severe milk-protein allergy or those who have hypersensitivity to umeclidinium, vilanterol or any excipients. **Side Effects:** Nasopharyngitis, oropharyngeal pain, sinusitis, pharyngitis, cough, urinary tract infection, constipation, dry mouth, hypertension, upper respiratory tract infections. **Warnings and Precautions:** Not recommended for use in patients with asthma or for relief of acute symptoms or an acute exacerbation. Use care when co-administering with strong CYP3A4 inhibitors (e.g. ketoconazole), beta-blockers and in patients with severe cardiovascular disease, narrow-angle glaucoma or urinary retention. **Pregnancy:** There are insufficient data from the use of umeclidinium/vilanterol in pregnant women. Anoro Ellipta should be used during pregnancy only if the expected benefit to the mother justifies the potential risk to the fetus. Before prescribing Anoro Ellipta, please review the data sheet for information on dosage, contraindications, precautions, interactions and adverse effects. The data sheet is available at [www.medsafe.govt.nz](http://www.medsafe.govt.nz).

Spiolto and Respimat are registered trade marks of Boehringer Ingelheim. Anoro and Ellipta are registered trade marks of the GlaxoSmithKline group of companies. The data sheets are available at [www.medsafe.govt.nz](http://www.medsafe.govt.nz). Anoro Ellipta was developed in collaboration with Innoviva Inc. Marketed by GlaxoSmithKline, NZ Limited, Auckland. **Adverse events involving GlaxoSmithKline products should be reported to GSK Medical Information on 0800 808 500. TAPS DA2123AM-PM-NZ-UCV-LBND-200004 Date of approval: Feb 2021.**

©2021 GSK group of companies or its licensor